

The Beckman-Quarles theorem for continuous mappings from $\mathbb{R}^2$ to $\mathbb{C}^2$

Apoloniusz Tyszk

IMUJ Preprint 2002/

Abstract

Let $\varphi : \mathbb{C}^2 \times \mathbb{C}^2 \rightarrow \mathbb{C}$, $\varphi((x_1, x_2), (y_1, y_2)) = (x_1 - y_1)^2 + (x_2 - y_2)^2$. We say that $f : \mathbb{R}^2 \rightarrow \mathbb{C}^2$ preserves distance $d \geq 0$ if for each $x, y \in \mathbb{R}^2$ $\varphi(x, y) = d^2$ implies $\varphi(f(x), f(y)) = d^2$. We prove that if $x, y \in \mathbb{R}^2$ and $|x - y| = (2\sqrt{2}/3)^k \cdot (\sqrt{3})^l$ (k, l are non-negative integers) then there exists a finite set $\{x, y\} \subseteq S_{xy} \subseteq \mathbb{R}^2$ such that each unit-distance preserving mapping from S_{xy} to $\mathbb{C}^2$ preserves the distance between x and y . It implies that each continuous map from $\mathbb{R}^2$ to $\mathbb{C}^2$ preserving unit distance preserves all distances.

The classical Beckman-Quarles theorem states that each unit-distance preserving mapping from $\mathbb{R}^n$ to $\mathbb{R}^n$ ($n \geq 2$) is an isometry, see [1], [2], [5] and [8]. Author's discrete form of this theorem ([9],[10]) states that if $x, y \in \mathbb{R}^n$ ($n \geq 2$) and $|x - y|$ is an algebraic number then there exists a finite set $\{x, y\} \subseteq S_{xy} \subseteq \mathbb{R}^n$ such that each unit-distance preserving mapping from S_{xy} to $\mathbb{R}^n$ preserves the distance between x and y .

Mathematics Subject Classification (2000): 51M05

Let $\varphi_n : \mathbb{C}^n \times \mathbb{C}^n \rightarrow \mathbb{C}$, $\varphi_n((x_1, \dots, x_n), (y_1, \dots, y_n)) = (x_1 - y_1)^2 + \dots + (x_n - y_n)^2$. We say that $f : \mathbb{R}^n \rightarrow \mathbb{C}^n$ preserves distance $d \geq 0$ if for each $x, y \in \mathbb{R}^n$ $\varphi_n(x, y) = d^2$ implies $\varphi_n(f(x), f(y)) = d^2$. In [11] the author proved that each continuous mapping from $\mathbb{R}^n$ to $\mathbb{C}^n$ ($n \geq 3$) preserving unit distance preserves all distances. In this paper we prove it for $n = 2$, similarly to the case $n \geq 3$ the proof is based on calculations using the Cayley-Menger determinant.

Proposition 1 ([11], cf. [3], [4]). The points $c_1 = (z_{1,1}, \dots, z_{1,n}), \dots, c_{n+1} = (z_{n+1,1}, \dots, z_{n+1,n}) \in \mathbb{C}^n$ are affinely dependent if and only if their Cayley-Menger determinant

$$\det \begin{bmatrix} 0 & 1 & 1 & \dots & 1 \\ 1 & \varphi_n(c_1, c_1) & \varphi_n(c_1, c_2) & \dots & \varphi_n(c_1, c_{n+1}) \\ 1 & \varphi_n(c_2, c_1) & \varphi_n(c_2, c_2) & \dots & \varphi_n(c_2, c_{n+1}) \\ \dots & \dots & \dots & \dots & \dots \\ 1 & \varphi_n(c_{n+1}, c_1) & \varphi_n(c_{n+1}, c_2) & \dots & \varphi_n(c_{n+1}, c_{n+1}) \end{bmatrix}$$

equals 0.

Proof. It follows from the equality

$$\left(\det \begin{bmatrix} z_{1,1} & z_{1,2} & \dots & z_{1,n} & 1 \\ z_{2,1} & z_{2,2} & \dots & z_{2,n} & 1 \\ \dots & \dots & \dots & \dots & \dots \\ z_{n+1,1} & z_{n+1,2} & \dots & z_{n+1,n} & 1 \end{bmatrix} \right)^2 =$$

$$\frac{(-1)^{n+1}}{2^n} \cdot \det \begin{bmatrix} 0 & 1 & 1 & \dots & 1 \\ 1 & \varphi_n(c_1, c_1) & \varphi_n(c_1, c_2) & \dots & \varphi_n(c_1, c_{n+1}) \\ 1 & \varphi_n(c_2, c_1) & \varphi_n(c_2, c_2) & \dots & \varphi_n(c_2, c_{n+1}) \\ \dots & \dots & \dots & \dots & \dots \\ 1 & \varphi_n(c_{n+1}, c_1) & \varphi_n(c_{n+1}, c_2) & \dots & \varphi_n(c_{n+1}, c_{n+1}) \end{bmatrix}.$$

Proposition 2 (see [11] for $k = 2$, cf. [3], [4]). For each points $c_1, \dots, c_{n+k} \in \mathbb{C}^n$ ($k = 2, 3, 4, \dots$) their Cayley-Menger determinant equals 0 i.e.

$$\det \begin{bmatrix} 0 & 1 & 1 & \dots & 1 \\ 1 & \varphi_n(c_1, c_1) & \varphi_n(c_1, c_2) & \dots & \varphi_n(c_1, c_{n+k}) \\ 1 & \varphi_n(c_2, c_1) & \varphi_n(c_2, c_2) & \dots & \varphi_n(c_2, c_{n+k}) \\ \dots & \dots & \dots & \dots & \dots \\ 1 & \varphi_n(c_{n+k}, c_1) & \varphi_n(c_{n+k}, c_2) & \dots & \varphi_n(c_{n+k}, c_{n+k}) \end{bmatrix} = 0.$$

Proof. Assume that $c_1 = (z_{1,1}, \dots, z_{1,n}), \dots, c_{n+k} = (z_{n+k,1}, \dots, z_{n+k,n})$. The points $\tilde{c}_1 = (z_{1,1}, \dots, z_{1,n}, 0, \dots, 0)$, $\tilde{c}_2 = (z_{2,1}, \dots, z_{2,n}, 0, \dots, 0)$, $\dots$, $\tilde{c}_{n+k} = (z_{n+k,1}, \dots, z_{n+k,n}, 0, \dots, 0) \in \mathbb{C}^{n+k-1}$ are affinely dependent. Since $\varphi_n(c_i, c_j) = \varphi_{n+k-1}(\tilde{c}_i, \tilde{c}_j)$ ($1 \leq i \leq j \leq n+k$) the Cayley-Menger determinant of points $c_1, \dots, c_{n+k}$ is equal to the Cayley-Menger determinant of points $\tilde{c}_1, \dots, \tilde{c}_{n+k}$ which equals 0 according to Proposition 1.

From Proposition 1 we obtain the following Propositions 3a and 3b.

Proposition 3a. If $c_1, c_2, c_3 \in \mathbb{C}^2$ and $\varphi_2(c_1, c_2) = \varphi_2(c_1, c_3) = \varphi_2(c_2, c_3) \in (0, \infty)$, then c_1, c_2, c_3 are affinely independent.

Proposition 3b. If $d > 0$, $c_1, c_2, c_3 \in \mathbb{C}^2$ and $\varphi_2(c_1, c_2) = 2d^2$, $\varphi_2(c_1, c_3) = 3d^2$, $\varphi_2(c_2, c_3) = 9d^2$, then c_1, c_2, c_3 are affinely independent.

Proposition 4 (see [11] for $\mathbb{C}^n$, cf. [4] p. 127 for $\mathbb{R}^n$). If $x, y, c_0, c_1, c_2 \in \mathbb{C}^2$, $\varphi_2(x, c_0) = \varphi_2(y, c_0)$, $\varphi_2(x, c_1) = \varphi_2(y, c_1)$, $\varphi_2(x, c_2) = \varphi_2(y, c_2)$ and c_0, c_1, c_2 are affinely independent, then $x = y$.

Proof. Computing we obtain that the vector $\overrightarrow{xy} := [s_1, s_2]$ is perpendicular to each of the linearly independent vectors $\overrightarrow{c_0c_1}$, $\overrightarrow{c_0c_2}$. Thus the vector $\overrightarrow{xy}$ is perpendicular to every linear combination of vectors $\overrightarrow{c_0c_1}$ and $\overrightarrow{c_0c_2}$. In particular, the vector $\overrightarrow{xy} = [s_1, s_2]$ is perpendicular to the vector $[\bar{s}_1, \bar{s}_2]$, where $\bar{s}_1, \bar{s}_2$ denote numbers conjugate to the numbers s_1, s_2 . Therefore $\overrightarrow{xy} = 0$ and the proof is complete.

From Kronecker's theorem (see [6]) we obtain the next proposition.

Proposition 5. The set $\{(2\sqrt{2}/3)^k \cdot (\sqrt{3})^l : k, l \in \{0, 1, 2, \dots\}\}$ is a dense subset of $(0, \infty)$.

Theorem 1. If $x, y \in \mathbb{R}^2$ and $|x - y| = (2\sqrt{2}/3)^k \cdot (\sqrt{3})^l$ (k, l are non-negative integers), then there exists a finite set $\{x, y\} \subseteq S_{xy} \subseteq \mathbb{R}^2$ such that each unit-distance preserving mapping from S_{xy} to $\mathbb{C}^2$ preserves the distance between x and y .

Proof. Let D denote the set of all positive numbers d with the following property:

- (*) if $x, y \in \mathbb{R}^2$ and $|x - y| = d$ then there exists a finite set $\{x, y\} \subseteq S_{xy} \subseteq \mathbb{R}^2$ such that any map $f : S_{xy} \rightarrow \mathbb{C}^2$ that preserves unit distance preserves also the distance between x and y .

Lemma 1 (see [11] for $n \geq 3$). If $d \in D$ then $\sqrt{3} \cdot d \in D$.

Proof. Let $d \in D$, $x, y \in \mathbb{R}^2$ and $|x - y| = \sqrt{3} \cdot d$. Using the notation of Figure 1 we show that

$$S_{xy} := S_{y\tilde{y}} \cup \bigcup_{i=1}^2 S_{xp_i} \cup \bigcup_{i=1}^2 S_{yp_i} \cup S_{p_1 p_2} \cup \bigcup_{i=1}^2 S_{x\tilde{p}_i} \cup \bigcup_{i=1}^2 S_{y\tilde{p}_i} \cup S_{\tilde{p}_1 \tilde{p}_2}$$

satisfies condition (*).




Figure 1

$$\begin{aligned} |x - y| &= |x - \tilde{y}| = \sqrt{3} \cdot d, |y - \tilde{y}| = |p_1 - p_2| = |\tilde{p}_1 - \tilde{p}_2| = d \\ |x - p_i| &= |y - p_i| = |x - \tilde{p}_i| = |\tilde{y} - \tilde{p}_i| = d \quad (i = 1, 2) \end{aligned}$$

Assume that $f : S_{xy} \rightarrow \mathbb{C}^2$ preserves unit distance. Since

$$S_{xy} \supseteq S_{y\tilde{y}} \cup \bigcup_{i=1}^2 S_{xp_i} \cup \bigcup_{i=1}^2 S_{yp_i} \cup S_{p_1 p_2}$$

we conclude that f preserves the distances between y and $\tilde{y}$, x and p_i ($i = 1, 2$), y and p_i ($i = 1, 2$), p_1 and p_2 . Hence $\varphi_2(f(y), f(\tilde{y})) = \varphi_2(f(x), f(p_i)) = \varphi_2(f(y), f(p_i)) = \varphi_2(f(p_1), f(p_2)) = d^2$ ($i = 1, 2$). By Proposition 2 the Cayley-Menger determinant of points $f(x)$, $f(p_1)$, $f(p_2)$, $f(y)$ equals 0 i.e.

$$\det \begin{bmatrix} 0 & 1 & 1 & 1 & 1 \\ 1 & \varphi_2(f(x), f(x)) & \varphi_2(f(x), f(p_1)) & \varphi_2(f(x), f(p_2)) & \varphi_2(f(x), f(y)) \\ 1 & \varphi_2(f(p_1), f(x)) & \varphi_2(f(p_1), f(p_1)) & \varphi_2(f(p_1), f(p_2)) & \varphi_2(f(p_1), f(y)) \\ 1 & \varphi_2(f(p_2), f(x)) & \varphi_2(f(p_2), f(p_1)) & \varphi_2(f(p_2), f(p_2)) & \varphi_2(f(p_2), f(y)) \\ 1 & \varphi_2(f(y), f(x)) & \varphi_2(f(y), f(p_1)) & \varphi_2(f(y), f(p_2)) & \varphi_2(f(y), f(y)) \end{bmatrix} = 0.$$

Denoting $t = \varphi_2(f(x), f(y))$ we obtain

$$\det \begin{bmatrix} 0 & 1 & 1 & 1 & 1 \\ 1 & 0 & d^2 & d^2 & t \\ 1 & d^2 & 0 & d^2 & d^2 \\ 1 & d^2 & d^2 & 0 & d^2 \\ 1 & t & d^2 & d^2 & 0 \end{bmatrix} = 0.$$

Computing this determinant we obtain

$$2d^2t \cdot (3d^2 - t) = 0.$$

Therefore

$$t = \varphi_2(f(x), f(y)) = \varphi_2(f(y), f(x)) = (\sqrt{3} \cdot d)^2$$

or

$$t = \varphi_2(f(x), f(y)) = \varphi_2(f(y), f(x)) = 0.$$

Analogously we may prove that

$$\varphi_2(f(x), f(\tilde{y})) = \varphi_2(f(\tilde{y}), f(x)) = (\sqrt{3} \cdot d)^2$$

or

$$\varphi_2(f(x), f(\tilde{y})) = \varphi_2(f(\tilde{y}), f(x)) = 0.$$

If $t = 0$ then the points $f(x)$ and $f(y)$ satisfy:

$$\begin{aligned} \varphi_2(f(x), f(x)) &= 0 = \varphi_2(f(y), f(x)), \\ \varphi_2(f(x), f(p_1)) &= d^2 = \varphi_2(f(y), f(p_1)), \\ \varphi_2(f(x), f(p_2)) &= d^2 = \varphi_2(f(y), f(p_2)). \end{aligned}$$

By Proposition 3a the points $f(x)$, $f(p_1)$, $f(p_2)$ are affinely independent. Therefore by Proposition 4 $f(x) = f(y)$ and consequently

$$d^2 = \varphi_2(f(y), f(\tilde{y})) = \varphi_2(f(x), f(\tilde{y})) \in \{(\sqrt{3} \cdot d)^2, 0\}.$$

Since $d^2 \neq (\sqrt{3} \cdot d)^2$ and $d^2 \neq 0$ we conclude that the case $t = 0$ cannot occur. This completes the proof of Lemma 1.

Lemma 2. If $d \in D$ then $3 \cdot d \in D$.

Proof. It follows from Lemma 1 because $3 \cdot d = \sqrt{3} \cdot (\sqrt{3} \cdot d)$.

Lemma 3. If $d \in D$ then $2 \cdot d \in D$.

Proof. Let $d \in D$, $x, y \in \mathbb{R}^2$ and $|x - y| = 2 \cdot d$. Using the notation of Figure 2 we show that

$S_{xy} := \bigcup \{S_{ab} : a, b \in \{x, y, p_1, p_2, p_3\}, |a - b| = d \vee |a - b| = \sqrt{3} \cdot d\}$
(where S_{xp_3} and S_{yp_2} are known to exist by Lemma 1) satisfies condition (*).




Figure 2

$$|x - y| = 2 \cdot d$$

$$|p_1 - p_2| = |p_1 - p_3| = |p_2 - p_3| = |x - p_1| = |x - p_2| = |y - p_1| = |y - p_3| = d$$

$$|x - p_3| = |y - p_2| = \sqrt{3} \cdot d$$

Assume that $f : S_{xy} \rightarrow \mathbb{C}^2$ preserves unit distance. Then f preserves all distances between p_i and p_j ($1 \leq i < j \leq 3$), x and p_i ($1 \leq i \leq 3$), y and p_i ($1 \leq i \leq 3$). By Proposition 2 the Cayley-Menger determinant of points $f(x), f(p_1), f(p_2), f(p_3), f(y)$ equals 0 i.e.

$$\det \begin{bmatrix} 0 & 1 & 1 & 1 & 1 & 1 \\ 1 & \varphi_2(f(x), f(x)) & \varphi_2(f(x), f(p_1)) & \varphi_2(f(x), f(p_2)) & \varphi_2(f(x), f(p_3)) & \varphi_2(f(x), f(y)) \\ 1 & \varphi_2(f(p_1), f(x)) & \varphi_2(f(p_1), f(p_1)) & \varphi_2(f(p_1), f(p_2)) & \varphi_2(f(p_1), f(p_3)) & \varphi_2(f(p_1), f(y)) \\ 1 & \varphi_2(f(p_2), f(x)) & \varphi_2(f(p_2), f(p_1)) & \varphi_2(f(p_2), f(p_2)) & \varphi_2(f(p_2), f(p_3)) & \varphi_2(f(p_2), f(y)) \\ 1 & \varphi_2(f(p_3), f(x)) & \varphi_2(f(p_3), f(p_1)) & \varphi_2(f(p_3), f(p_2)) & \varphi_2(f(p_3), f(p_3)) & \varphi_2(f(p_3), f(y)) \\ 1 & \varphi_2(f(y), f(x)) & \varphi_2(f(y), f(p_1)) & \varphi_2(f(y), f(p_2)) & \varphi_2(f(y), f(p_3)) & \varphi_2(f(y), f(y)) \end{bmatrix} = 0.$$

Denoting $t = \varphi_2(f(x), f(y))$ we obtain

$$\det \begin{bmatrix} 0 & 1 & 1 & 1 & 1 & 1 \\ 1 & 0 & d^2 & d^2 & 3d^2 & t \\ 1 & d^2 & 0 & d^2 & d^2 & d^2 \\ 1 & d^2 & d^2 & 0 & d^2 & 3d^2 \\ 1 & 3d^2 & d^2 & d^2 & 0 & d^2 \\ 1 & t & d^2 & 3d^2 & d^2 & 0 \end{bmatrix} = 0.$$

Computing this determinant we obtain

$$3d^4 \cdot (t - 4d^2)^2 = 0.$$

Therefore

$$t = \varphi_2(f(x), f(y)) = \varphi_2(f(y), f(x)) = (2d)^2.$$

Lemma 4. If $a, b \in D$ and $a > b$, then $\sqrt{a^2 - b^2} \in D$.

Proof. Let $a, b \in D$, $a > b$, $x, y \in \mathbb{R}^2$ and $|x - y| = \sqrt{a^2 - b^2}$. Using the notation of Figure 3 we show that

$$S_{xy} := S_{xp_1} \cup S_{xp_2} \cup S_{yp_1} \cup S_{yp_2} \cup S_{p_1p_2}$$

(where $S_{p_1p_2}$ is known to exist by Lemma 3) satisfies condition (*).




Figure 3

$$|x - y| = \sqrt{a^2 - b^2}$$

$$|x - p_1| = |x - p_2| = b, |y - p_1| = |y - p_2| = a, |p_1 - p_2| = 2b$$

Assume that $f : S_{xy} \rightarrow \mathbb{C}^2$ preserves unit distance. Then f preserves the distances between x and p_i ($i = 1, 2$), y and p_i ($i = 1, 2$), p_1 and p_2 . By Proposition 2 the Cayley-Menger determinant of points $f(x)$, $f(p_1)$, $f(p_2)$, $f(y)$ equals 0 i.e.

$$\det \begin{bmatrix} 0 & 1 & 1 & 1 & 1 \\ 1 & \varphi_2(f(x), f(x)) & \varphi_2(f(x), f(p_1)) & \varphi_2(f(x), f(p_2)) & \varphi_2(f(x), f(y)) \\ 1 & \varphi_2(f(p_1), f(x)) & \varphi_2(f(p_1), f(p_1)) & \varphi_2(f(p_1), f(p_2)) & \varphi_2(f(p_1), f(y)) \\ 1 & \varphi_2(f(p_2), f(x)) & \varphi_2(f(p_2), f(p_1)) & \varphi_2(f(p_2), f(p_2)) & \varphi_2(f(p_2), f(y)) \\ 1 & \varphi_2(f(y), f(x)) & \varphi_2(f(y), f(p_1)) & \varphi_2(f(y), f(p_2)) & \varphi_2(f(y), f(y)) \end{bmatrix} = 0.$$

Denoting $t = \varphi_2(f(x), f(y))$ we obtain

$$\det \begin{bmatrix} 0 & 1 & 1 & 1 & 1 \\ 1 & 0 & b^2 & b^2 & t \\ 1 & b^2 & 0 & 4b^2 & a^2 \\ 1 & b^2 & 4b^2 & 0 & a^2 \\ 1 & t & a^2 & a^2 & 0 \end{bmatrix} = 0.$$

Computing this determinant we obtain

$$-8b^2 \cdot (t + b^2 - a^2)^2 = 0.$$

Therefore

$$t = a^2 - b^2.$$

Lemma 5. If $d \in D$ then $\sqrt{2} \cdot d \in D$.

Proof. It follows from equality $\sqrt{2} \cdot d = \sqrt{(\sqrt{3} \cdot d)^2 - d^2}$ and Lemmas 1 and 4.

Lemma 6. If $d \in D$ then $(2\sqrt{2}/3) \cdot d \in D$.

Proof. Let $d \in D$, $x, y \in \mathbb{R}^2$, $|x - y| = (2\sqrt{2}/3) \cdot d$. Using the notation of Figure 4 we show that

$$S_{xy} := S_{y\tilde{y}} \cup \bigcup_{i=1}^2 S_{xp_i} \cup \bigcup_{i=1}^2 S_{yp_i} \cup S_{p_1p_2} \cup \bigcup_{i=1}^2 S_{x\tilde{p}_i} \cup \bigcup_{i=1}^2 S_{y\tilde{p}_i} \cup S_{\tilde{p}_1\tilde{p}_2}$$

(where sets corresponding to distances $\sqrt{3} \cdot d$ are known to exist by Lemma 1, sets corresponding to distances $\sqrt{2} \cdot d$ are known to exist by Lemma 5, sets corresponding to distances $3 \cdot d$ are known to exist by Lemma 2) satisfies condition (*).




Figure 4

$$\begin{aligned}
|x - y| &= |x - \tilde{y}| = (2\sqrt{2}/3) \cdot d \\
|y - \tilde{y}| &= d \\
|x - p_1| &= |y - p_1| = |x - \tilde{p}_1| = |\tilde{y} - \tilde{p}_1| = \sqrt{3} \cdot d \\
|x - p_2| &= |y - p_2| = |x - \tilde{p}_2| = |\tilde{y} - \tilde{p}_2| = \sqrt{2} \cdot d \\
|p_1 - p_2| &= |\tilde{p}_1 - \tilde{p}_2| = 3 \cdot d
\end{aligned}$$

Assume that $f : S_{xy} \rightarrow \mathbb{C}^2$ preserves unit distance. Since

$$S_{xy} \supseteq S_{y\tilde{y}} \cup \bigcup_{i=1}^2 S_{xp_i} \cup \bigcup_{i=1}^2 S_{yp_i} \cup S_{p_1p_2}$$

we conclude that f preserves the distances between y and $\tilde{y}$, x and p_i ($i = 1, 2$), y and p_i ($i = 1, 2$), p_1 and p_2 . By Proposition 2 the Cayley-Menger determinant of points $f(x)$, $f(p_1)$, $f(p_2)$, $f(y)$ equals 0 i.e.

$$\det \begin{bmatrix} 0 & 1 & 1 & 1 & 1 \\ 1 & \varphi_2(f(x), f(x)) & \varphi_2(f(x), f(p_1)) & \varphi_2(f(x), f(p_2)) & \varphi_2(f(x), f(y)) \\ 1 & \varphi_2(f(p_1), f(x)) & \varphi_2(f(p_1), f(p_1)) & \varphi_2(f(p_1), f(p_2)) & \varphi_2(f(p_1), f(y)) \\ 1 & \varphi_2(f(p_2), f(x)) & \varphi_2(f(p_2), f(p_1)) & \varphi_2(f(p_2), f(p_2)) & \varphi_2(f(p_2), f(y)) \\ 1 & \varphi_2(f(y), f(x)) & \varphi_2(f(y), f(p_1)) & \varphi_2(f(y), f(p_2)) & \varphi_2(f(y), f(y)) \end{bmatrix} = 0.$$

Denoting $t = \varphi_2(f(x), f(y))$ we obtain

$$\det \begin{bmatrix} 0 & 1 & 1 & 1 & 1 \\ 1 & 0 & 3d^2 & 2d^2 & t \\ 1 & 3d^2 & 0 & 9d^2 & 3d^2 \\ 1 & 2d^2 & 9d^2 & 0 & 2d^2 \\ 1 & t & 3d^2 & 2d^2 & 0 \end{bmatrix} = 0.$$

Computing this determinant we obtain

$$2d^2t \cdot (8d^2 - 9t) = 0.$$

Therefore

$$t = \varphi_2(f(x), f(y)) = \varphi_2(f(y), f(x)) = ((2\sqrt{2}/3) \cdot d)^2$$

or

$$t = \varphi_2(f(x), f(y)) = \varphi_2(f(y), f(x)) = 0.$$

Analogously we may prove that

$$\varphi_2(f(x), f(\tilde{y})) = \varphi_2(f(\tilde{y}), f(x)) = ((2\sqrt{2}/3) \cdot d)^2$$

or

$$\varphi_2(f(x), f(\tilde{y})) = \varphi_2(f(\tilde{y}), f(x)) = 0.$$

If $t = 0$ then the points $f(x)$ and $f(y)$ satisfy:

$$\begin{aligned} \varphi_2(f(x), f(x)) &= 0 = \varphi_2(f(y), f(x)), \\ \varphi_2(f(x), f(p_1)) &= 3d^2 = \varphi_2(f(y), f(p_1)), \\ \varphi_2(f(x), f(p_2)) &= 2d^2 = \varphi_2(f(y), f(p_2)). \end{aligned}$$

By Proposition 3b the points $f(x), f(p_1), f(p_2)$ are affinely independent. Therefore by Proposition 4 $f(x) = f(y)$ and consequently

$$d^2 = \varphi_2(f(y), f(\tilde{y})) = \varphi_2(f(x), f(\tilde{y})) \in \{((2\sqrt{2}/3) \cdot d)^2, 0\}.$$

Since $d^2 \neq ((2\sqrt{2}/3) \cdot d)^2$ and $d^2 \neq 0$ we conclude that the case $t = 0$ cannot occur. This completes the proof of Lemma 6.

Obviously $1 \in D$. Therefore by Lemmas 1 and 6

$$\{(2\sqrt{2}/3)^k \cdot (\sqrt{3})^l : k, l \in \{0, 1, 2, \dots\}\} \subseteq D.$$

This completes the proof of Theorem 1.

As a corollary of Theorem 1 and Proposition 5 we obtain our main theorem.

Theorem 2. Each continuous map from $\mathbb{R}^2$ to $\mathbb{C}^2$ preserving unit distance preserves all distances.

Remark ([11]). By an endomorphism of $\mathbb{C}$ we understand any map $f : \mathbb{C} \rightarrow \mathbb{C}$ satisfying:

$$\begin{aligned}\forall x, y \in \mathbb{C} \quad f(x + y) &= f(x) + f(y), \\ \forall x, y \in \mathbb{C} \quad f(x \cdot y) &= f(x) \cdot f(y), \\ f(0) &= 0, \\ f(1) &= 1.\end{aligned}$$

If $f : \mathbb{C} \rightarrow \mathbb{C}$ is an endomorphism then $(f|_{\mathbb{R}}, \dots, f|_{\mathbb{R}}) : \mathbb{R}^n \rightarrow \mathbb{C}^n$ preserves unit distance. Bijective endomorphisms are called automorphisms. There are two trivial automorphisms of $\mathbb{C}$: identity and conjugation. It is known that there exist non-trivial automorphisms of $\mathbb{C}$ and each such automorphism $f : \mathbb{C} \rightarrow \mathbb{C}$ satisfies: $\exists \frac{x \in \mathbb{R}}{x \neq 0} f(x) \notin \mathbb{R}$ ([7]). From this $\varphi_n((0, 0, \dots, 0), (x, 0, \dots, 0)) = |x|^2$ and $\varphi_n((f(0), f(0), \dots, f(0)), (f(x), f(0), \dots, f(0))) = (f(x))^2 \neq |x|^2$. Therefore $(f|_{\mathbb{R}}, \dots, f|_{\mathbb{R}})$ preserves unit distance, but does not preserve the distance $|x| > 0$.

References

- [1] F. S. Beckman and D. A. Quarles Jr., *On isometries of euclidean spaces*, Proc. Amer. Math. Soc. 4 (1953), 810–815.
- [2] W. Benz, *Geometrische Transformationen (unter besonderer Berücksichtigung der Lorentztransformationen)*, BI Wissenschaftsverlag, Mannheim, 1992.
- [3] L. M. Blumenthal, *Theory and applications of distance geometry*, 2nd ed., Chelsea Publishing Co., New York, 1970.
- [4] K. Borsuk, *Multidimensional analytic geometry*, Polish Scientific Publishers, Warsaw, 1969.
- [5] U. Everling, *Solution of the isometry problem stated by K. Ciesielski*, Math. Intelligencer 10 (1988), No. 4, p. 47.

- [6] G. H. Hardy and E. M. Wright, *An introduction to the theory of numbers*, 5th ed., The Clarendon Press, Oxford University Press, New York, 1979.
- [7] M. Kuczma, *An introduction to the theory of functional equations and inequalities: Cauchy's equation and Jensen's inequality*, Polish Scientific Publishers and Silesian University, Warsaw-Cracow-Katowice, 1985.
- [8] J. A. Lester, *Distance preserving transformations*, in: "Handbook of incidence geometry" (ed. F. Buekenhout), 921–944, North-Holland, Amsterdam, 1995.
- [9] A. Tyszką, *Discrete versions of the Beckman-Quarles theorem*, Aequationes Math. 59 (2000), 124–133.
- [10] A. Tyszką, *Discrete versions of the Beckman-Quarles theorem from the definability results of R. M. Robinson*, Algebra, Geometry & their Applications, Seminar Proceedings 1 (2001), 88–90, Yerevan State University Press.
- [11] A. Tyszką, *The Beckman-Quarles theorem for continuous mappings from $\mathbb{R}^n$ to $\mathbb{C}^n$* , IMUJ Preprint 2002/07, Instytut Matematyki, Uniwersytet Jagielloński, Kraków, 2002.

Apoloniusz Tyszką
 Technical Faculty
 Hugo Kołłątaj University
 Balicka 104
 30-149 Kraków, Poland
 E-mail: rttyszka@cyf-kr.edu.pl